

First isolation of West Nile virus in Brazil

Lívia Carício Martins¹, Eliana Vieira Pinto da Silva¹, Livia Medeiros Neves Casseb¹, Sandro Patroca da Silva¹, Ana Cecília Ribeiro Cruz¹, Jamilla Augusta de Sousa Pantoja¹, Daniele Barbosa de Almeida Medeiros¹, Arnaldo Jorge Martins Filho², Ermelinda do Rosário Moutinho da Cruz², Marialva Tereza Ferreira de Araújo², Jedson Ferreira Cardoso³, Marcos Antônio Correia Rodrigues da Cunha⁴, Gilton Luiz Almada⁴, Alessandro Pecego Martins Romano⁵, Maria Guadalupe Dias Pestana Santos⁶, Gilsa Aparecida Pimenta Rodrigues⁴, Jannifer Oliveira Chiang¹, Juarez Antonio Simões Quaresma², Valéria Lima Carvalho^{1/+}, Pedro Fernando da Costa Vasconcelos¹

¹Instituto Evandro Chagas, Seção de Arbovirologia e Febres Hemorrágicas, Ananindeua, PA, Brasil

²Instituto Evandro Chagas, Seção de Patologia, Ananindeua, PA, Brasil

³Instituto Evandro Chagas, Centro de Inovação tecnológica, Ananindeua, PA, Brasil

⁴Secretaria de Estado da Saúde do Espírito Santo, Vitória, ES, Brasil

⁵Ministério da Saúde, Secretaria de Vigilância em Saúde, Brasília, DF, Brasil

⁶Secretaria de Saúde do Município de Venécia, Venécia, ES, Brasil

BACKGROUND Serological evidence of West Nile virus (WNV) infection has been reported in different regions of Brazil from equine and human hosts but the virus had never been isolated in the country.

OBJECTIVES We sought to identify the viral etiology of equine encephalitis in Espírito Santo state.

METHODS We performed viral culture in C6/36 cells, molecular detection of WNV genome, histopathology and immunohistochemistry from horse cerebral tissue. We also carried out sequencing, phylogenetic analysis and molecular clock.

FINDINGS Histopathologic analysis from horse cerebral tissue showed injury related to encephalitis and WNV infection was confirmed by immunohistochemistry. The virus was detected by reverse transcription quantitative polymerase chain reaction (RT-qPCR) from brain tissue and subsequently isolated in C6/36 cells. WNV full-length genome was sequenced showing the isolated strain belongs to lineage 1a. The molecular clock indicated that Brazilian WNV strain share the same common ancestor that were circulating in US during 2002-2005.

MAIN CONCLUSIONS Here we report the first isolation of WNV in Brazil from a horse with neurologic disease, which was clustered into lineage 1a with others US WNV strains isolated in beginning of 2000's decade.

Key words: West Nile virus - horses - Brazil

West Nile virus (WNV) is a member of the Japanese Encephalitis serocomplex within *Flaviviridae* family and *Flavivirus* genus.⁽¹⁾ Originally, WNV was isolated from a human with fever in Uganda, 1937.⁽²⁾ Later it would be detected in the Middle East, France, South Asia and Australia. In 1999, the virus was introduced into USA and since then, it has spread from Canada to Argentina.⁽³⁾

WNV is maintained in nature by transmission between *Culex* mosquitoes, in particular *Cx. pipiens*, considered its principal vector, and certain passerine birds, its main reservoir host.⁽⁴⁾ Horses and humans are incidental or "dead-end" hosts, since they develop low level viremia which is insufficient to contribute to WNV spread via mosquitoes.^(4,5)

WNV is a neuropathogen to human, equine, and many other mammalian and avian species.⁽⁴⁾ In humans, around 80% of infected people are asymptomatic and 20% will develop West Nile Fever (WNF) which can range from mild acute febrile illness, to neurological diseases including meningitis, encephalitis, and acute flaccid paralysis.⁽⁶⁾ Regarding horses, when they develop disease the main symptoms are fever, depression, loss of appetite, colic, encephalitis with ataxia, limb weakness, recumbency and muscle fasciculation.⁽⁷⁾

Phylogenetic analysis established nine WNV lineages according to geographic location and the type of host. Lineages 1 and 2 are associated with the majority of human outbreaks in Europe, Africa, Middle East, Americas and Oceania. Lineages 3, 4, 6, and 9 are from Europe. Lineage 5 is from India. Lineages 7 and 8 are from Africa.⁽¹⁾

In South America, WNV was isolated for the first time in Argentina from horses in 2006 and two years later, the virus was isolated in Colombia from captive flamingoes.^(8,9) Regarding Brazil, after the spread of the virus in North America, in 2003, the National WNV Surveillance System was created, based on the recommendations of Pan American Health Organization (PAHO) and World Health Organization (WHO). Since then, any

doi: 10.1590/0074-02760180332

Financial support: Instituto Evandro Chagas, Ministério da Saúde. PFCV (grants 303999/2016-0 and 4399771/2016-0) and DBAM (grant 306581/2016-7) were partially supported by CNPq.

+ Corresponding author: valeriacarvalho@iec.gov.br

http://orcid.org/0000-0001-8385-8253

Received 13 July 2018

Accepted 27 December 2018



suspected human or animal case of WNF is investigated following the established protocol and biological samples are referred for differential laboratory diagnosis.

Over the years, some studies reported serological evidence of WNV in Brazil from different hosts and regions indicating the circulation of the virus,^(10,11,12,13) but the virus had never been isolated in the country. We describe the first isolation of WNV in Brazil, obtained from a horse with encephalitis in Espírito Santo state.

MATERIALS AND METHODS

Clinical information - An adult male horse from Pedra Grande region on the São Mateus municipality, Espírito Santo state, Brazil, with signs of colic developed neurological manifestations on the 25th of April 2018 including dysphagia, ataxia in anterior limbs, muscle tremors, shaking, opisthotonos, lateral decubitus, intense sweating, pedaling movements and other alterations indicating hemineglect. Within 24 hours, the animal showed signs of paralysis of the pelvic limbs, difficulty chewing, and not responding to needle stimulation along the spine. The animal was euthanised and specimens were submitted to the Evandro Chagas Institute, National Reference Laboratory for Arbovirus, for laboratory diagnosis, including testing for WNV. The sample was assigned registration number BE AN 854747.

Before the disease episode, the animal had been immunised with Tri-Equi-Hertape vaccine covering protection against Eastern equine encephalitis virus, Western equine encephalitis virus, equine influenza viruses (types A1 and A2, including Kentucky 92) and tetanus toxoid.

Virus isolation - Approximately 0.05 gram of central nervous system (CNS) sample was homogenised in 1.0 mL phosphate-buffered saline, pH 7.4, containing 5% fetal bovine serum with penicillin (100 U/mL), streptomycin (100 mg/mL) and gentamicin (0.05 mg/mL), using one 5 mm stainless steel bead in a TissueLyser (Qiagen, Hilden, Germany) set to 26 Hz for four min. The sample was frozen at -80°C overnight, then thawed and centrifuged at 16,266 g for 10 min (4°C) and then further clarified with a 0.22-µm nylon syringe filter. Then 100 µL of the supernatant filtrate was inoculated into C6/36 cells and after 1 h of adsorption at 28°C, 1.5 mL of Leibovitz's L-15 maintenance medium (supplemented with 2% Foetal Bovine Serum, 0.1% Penicillin 100 U/mL and Streptomycin 100 mg/mL, 10% Tryptose Phosphate Broth and 1% Non-essential amino acids) was added into 20 cm² tissue culture tube. The cells were incubated at 28°C and observed during seven days, scraped from the tubes and collected on slides to perform indirect immunofluorescence assay (IFA) using hyperimmune mouse ascitic fluids to *Flavivirus* group-reactive (including antibodies to the following viruses: Bussuquara, Ilhéus, Rocio, Cacipacoré, Dengue 1, 2, 3, 4, Naranjal, Zika and West Nile-chimera), WNV, as well as, to *Alphavirus* group-reactive and Oropouche virus to rule out infection by other families of viruses.

Histopathology and immunohistochemistry - Paraffin-embedded tissue samples were processed for histopathology and stained with hematoxylin and eosin (HE).

⁽¹⁴⁾ For immunohistochemistry, an adapted Streptavidin Alkaline Phosphatase assay⁽¹⁵⁾ with anti-WNV polyclonal antibody serum was used.

Detection of WNV genome - The RNA was extracted from CNS tissues supernatant by using Maxwell 16 Tissue LEV Total RNA Purification Kit (Promega) using Maxwell automated system. Reverse transcription quantitative polymerase chain reaction (RT-qPCR) for detection of WNV genome was performed following established protocol.⁽¹⁶⁾ The assay was performed on a 7500 Real Time PCR System (Applied Biosystems) using Superscript III Platinum One-Step qRT-PCR System kit (Invitrogen). We also performed RT-qPCR for detection of Saint Louis encephalitis virus, for differential laboratory diagnosis, as described.⁽¹⁷⁾

Sequencing - The cell culture supernatant was extracted with QIAamp Viral RNA Mini Kit. The metagenome sequencing process started with the ssRNA. The synthesis of the first and second strand was performed using SuperScript VILO MasterMix and NEBNext Second Strand Synthesis Module, respectively. The reaction was purified with PureLink PCR Purification Kit, following manufacturer's instructions.

The cDNA library was prepared and sequencing using the methodology described in the Nextera XT DNA Library Preparation Kit on a Miniseq (Illumina, Inc).

The genome sequence was determined using the *De Novo Assembler* methodology in IDBA-UD program⁽¹⁸⁾ and SPAdes.⁽¹⁹⁾ The inspection, annotations of putative open reading frames (ORF) genes and additional analysis were performed using the Geneious v.9.1.6 software (Biomatters, New Zealand). All contigs were aligned and compared against the database of virus proteins available in NCBI through the Diamond.⁽²⁰⁾

Phylogenetic and evolutionary analysis - Initially, a multiple sequencing alignment, using the entire ORFs of the Brazilian strain and more than 1,500 WNV nucleotide sequences available on NCBI, was performed using Mafft v7.310 software⁽²¹⁾ and for visual inspection we used Geneious v.9.1.8 software. Subsequently, recombination events were evaluated using the Phi-test implemented on SplitsTree4 v4.14.6 program.⁽²²⁾ The best-fitting model of nucleotide substitution was determined using jModelTest v.2.1.10.⁽²³⁾ A maximum likelihood (ML) tree was constructed using the RAxML v.8.2.11 software.⁽²⁴⁾ Afterward, a new nucleotide dataset was selected to perform evolutionary analysis based on the full phylogenetic tree using the entire ORFs of 166 WNV strains comprising other South American WNV strains and basal lineages to the cluster including the Brazilian WNV strain.

The temporal structures for each one and the combined WNV clades were verified by using the TempEst v1.5.1, and all data were plotted using ggplot2 with the R programming language. To estimate the time to the most recent common ancestor (MRCA) for each monophyletic clusters and the rate of nucleotide substitutions per site, we used the Markov Chain Monte Carlo (MCMC) employing Bayesian approach available on the BEAUTI and BEAST v.1.10.1 package and BEAGLE library. However, we firstly established the prior MCMC chosen by

testing all models and determining Bayes factors ($\log_{10} BF > 3$), as well as, the best clock and coalescent models were selected using the log marginal likelihood estimation (MLE) under path sampling (PS) and stepping-stone sampling (SS) methods implemented in BEAST. The best model used was the general time-reversible with invariant sites (GTR + I + T4) (Supplementary data II).




Fig. 1: detection of West Nile virus (WNV) antigen in C6/36 cells by indirect immunofluorescence assay (IFA). (A) WNV antigen (green) in infected cells detected by IFA using flavivirus group-reactive hyperimmune mouse ascitic fluid. (B) Uninfected cell control by IFA using the same hyperimmune mouse ascitic fluid.

We calculated different coalescent models based parametric Constant and non-parametric was the Skygrid and Skyline models, as well as, the molecular clock models using the strict molecular clock (Strict) and the uncorrelated relaxed lognormal molecular clock (UCLN). The MCMC method was employed with three independently runs each composed by 100 million generations and uncertainties of the parameters estimates were excluded after assessing the initial 10% of burn-in by calculating the Effective Sample Size (ESS) within a confidence interval of 95% Highest Probability Density (HPD) value, using the TRACER v.1.7.1.⁽²⁵⁾ A maximum clade credibility tree (MCC) was produced and annotated by the use of Tree-Annotator in the BEAST using FigTree v.1.4.3.⁽²⁶⁾

RESULTS

Virus isolation and detection of WNV genome - C6/36 cells inoculated with the sample BE AN 854747 (CNS material) presented cytopathic effect (CPE) in the fourth day post-inoculation (dpi) characterised by cell death and formation of syncytia. On the seventh dpi, the indirect immunofluorescence assay showed approximately 75% positive cells for both *Flavivirus* and WNV polyclonal antibodies (Fig. 1). The horse's brain samples were also positive to WNV by RT-qPCR (Ct: 31.2).

Histopathology and immunohistochemistry - The histological sections of the horse's brain showed tissue alterations characterised by inflammatory infiltration of the parenchyma, which is sometimes distributed around the vessels and consist of mononuclear and polymorphonuclear leukocytes, area of variable neuronal necrosis, neuronophagia, microglial nodules and gliosis. Anti-WNV




Fig. 2: photomicrograph of cerebral tissue of a horse with neurologic manifestation from Espírito Santo state with description of histopathological aspects and positive immunostaining for West Nile virus (WNV). (A) Inflammatory infiltrate predominantly constituted by mononuclear cells with perivascular cuffing (arrow) and neuronal necrosis (arrow heads). (B) Brain tissue showed neuronophagy (arrows) and associated gliosis (arrow head). (C) Perivascular infiltrate with congestion (arrows) and foci of neuronal necrosis of different intensities (arrow heads). (D) Immunohistochemistry of WNV antigen (reddish staining) in the cell cytoplasm in several foci of brain tissue (arrows).




Fig. 3: midpoint phylogenetic tree of nucleotide sequences using only polypeptide coding region of 1582 West Nile virus (WNV) strains representing different viral lineages. The analysis of those nucleotide sequences was performed using the maximum likelihood method based on the GTR matrix-based model. Different phylogenetic groups are assigned to previously defined lineages; highlighted in dark blue, strains included in lineage 1a from USA and in light blue strains included in lineage 1a from Europe, Asia and Africa. Strains from Brazil (red), Mexico (orange), Colombia (purple), and Argentina (green) were also identified on the phylogenetic tree. Numbers over each main node of the tree correspond to bootstrap values (1000 replicates). Values < 70 are not supported by reproducible topologies. The bar represents nucleotide substitutions along the branch.

antibodies represented by deposition of reddish granules in the cytoplasm of the cells showed the presence of viral antigens in the cerebral parenchyma (Fig. 2).


Phylogenetic and evolutionary analysis - The ML phylogenetic trees, using GTR+I+T4 model, showed similar topology identifying all WNV lineages previously described. Our data showed that the BE AN 854747 strain (GenBank accession number: MH643887) belonged to lineage 1a related to others strains from United States and Mexico (Fig. 3; Supplementary data I). The Brazilian strain was more closely related with two US strains isolated from *Corvus brachyrhynchos* (KJ501489) and *Pelecanus erythrorhynchos* (KJ5011492) in 2002 and 2005, respectively, with average of identity of 99.0% (nucleotide) and 99.6% (amino acid).

To build molecular clock tree, we checked recombination to lineages 1a and 1b. The Phi-test did not find statistical significant evidence for recombination ($p = 0.1979$). Furthermore, 166 WNV strain, including the Brazilian strain were previously selected and a linear regression analysis was performed (Fig. 4A) based on ML tree (Supplementary data II), showing good temporal signal for this dataset.

The best molecular clock and coalescent model were uncorrelated relaxed lognormal molecular clock (UCLN) and Skyline models, respectively (Fig. 4B). The evolutionary analysis of the 166 WNV strains isolated from 1960 to 2018 indicates that the most recent common ancestor of all South America strains were from US and probably were introduced by independent events in Argentina, Brazil and Colombia where the estimated time were 16.5 years ago (15.79 - 17.39, 95% HPD), 17.2 years ago (16.34 - 18.06, 95% HPD) and 11.2 years ago (10.64 - 11.76, 95% HPD), suggesting that the ancestors of these clusters were circulating in 2002, 2000 and 2007, respectively. The introduction of Brazilian strain was estimated by most closely related sequences of USA KJ501492/2005 and KJ501489/2002 (Fig. 4C).

DISCUSSION

We report the first isolate and detection of WNV in Brazil from an equine CNS sample collected in a rural area of Espírito Santo state. The virus caused histopathologic alterations in the cerebral tissue of the horse compatible with encephalitis and similar to those previously described for WNV.⁽⁶⁾ Infected C6/36 cells presented char-



acteristic morphologic changes commonly observed for flaviviruses. The phylogenetic analysis of the full-length genome showed that this WNV isolate is included in lineage 1a, which comprises strains isolated from Europe, Africa, and the Americas.⁽¹⁾ We also identified greater genetic relatedness of the BE AN 854747 isolate with North American strains as well as Central (Mexico) and South America strains (Colombia and Argentina).^(27,8,9)

The molecular clock suggested that Brazilian WNV share the same ancestor with US strains that were circulating between 2000 to 2005 and probably was introduced in Brazil from a different event than Argentine and Colombian. We observed that Mexico presented multiple introductions all of them related to US strains and the oldest Mexican strains, introduced in 2001, are most closely to the Brazilian strain (BE AN 854747).

Since 1999 when WNV emerged in USA, the virus has spread throughout the Americas from Canada to Argentina causing several thousand cases of neurological disease, including cases of fatalities in humans, but with higher rates of mortality in birds and horses. After the detection of the virus in the Americas, many efforts have been made to detect the possible circulation of the virus in Brazil. Thus, neutralising antibodies were first detected in equines and chickens in Pantanal region in 2009 and 2010;^(10,11) and later in equines in the Northeast region and again in the Pantanal.⁽¹²⁾ In 2015, the country reported the first human case of WNV encephalitis with flaccid paralysis in Piauí state in which all CDC confirmation criteria for WNF were filled including the detection of neutralising antibodies against the virus.⁽¹³⁾ This indicates that WNV had been previously enzootically circulating but never had been isolated in the country. It is worth pointing out that in Brazil there is the co-circulation of other flaviviruses like Dengue, Zika, Yellow Fever, Saint Louis encephalitis, Ilheus and others, which complicates the serological diagnosis of these viruses, due to the extensive flavivirus cross-reactivity in serological assays.⁽²⁸⁾ Furthermore, this cross-reactivity can lead to cross-protection which may prevent WNV from causing large epidemics in humans in Brazil. However, WNV may become an important pathogen to animals, especially wild life and animals of production, affecting directly the economy of the country. Currently the health, agricultural and environmental authorities are investigating this equine case and implications for local human and animal populations. At the national level, the Brazilian surveillance system investigates neuroinvasive arboviral disease, creating notification systems for both human and animal surveillance, as well as a guide to clinical management and establishment of sentinel hospitals to investigate neurologic cases in each state of the country.⁽²⁹⁾ Nevertheless, it is important that with the first isolation of WNV, the Brazilian surveillance systems be further alert to human and equine neurological cases compatible with WN Fever. The notification and investigation of neurological diseases in equidae as well as the epizootic events of wild birds should be improved as they may be the key to the detection of WNV before outbreaks of large magnitude.

Ethics - Biological samples of animals were obtained and sent by Epidemiologic Surveillance, Ministry of Health, to Evandro Chagas Institute, national reference laboratory, to confirm the laboratory diagnosis. The euthanasia of the animals was made by *anesthetic* overdose following parameters established by Guia para la Vigilancia, Detección y Respuesta para las Encefalitis Equinas.⁽³⁰⁾ This procedure was carried out for diagnostic purposes, and therefore does not require consideration by ethics committees according to Brazilian law number 11.794 of October 8th, 2008.

ACKNOWLEDGEMENTS

To Maria de Lourdes Gomes Lima for her help with the immunohistochemistry. Our acknowledgments also to the professionals of Secretariat of Health and Agricultural Defense of Espírito Santo state and the horse's owner that allowed and supported the euthanasia and sample collection. We also would like to thank Alice Louize Nunes Queiroz and Daniela Sueli Guerreiro Rodrigues for their help with the genome detection. Our acknowledgments to Mario Basara, a native English speaker, for the English revision.

AUTHORS' CONTRIBUTION

LCM, PFCV and JOC - Supervision; VLC, EVPS and ACRC - writing, original draft; APMR, LMNC, MACRC, GLA, MGDPS and GAPR - investigation; SPS, JASP, AJMF, ERM, MTF and JASQ - methodology; JFC and DBAM - molecular clock. The author declares no conflicts of interest.

REFERENCES

1. Fall G, Di Paola N, Faye M, Dia M, Freire CCM, Loucoubar C, et al. Biological and phylogenetic characteristics of West African lineages of West Nile virus (DWC Beasley, Ed.). *PLoS Negl Trop Dis*. 2017; 11(11): 1-23.
2. Smithburn KC, Hughes TP, Burke AW, Paul JH. A neurotropic virus isolated from the blood of a native of Uganda. *Am J Trop Med Hyg*. 1940; 20: 471-2.
3. Komar N, Clark GG. West Nile virus activity in Latin America and the Caribbean. *Rev Panam Salud Publica*. 2006; 19(2): 112-7.
4. Campbell GL, Marfin AA, Lanciotti RS, Gubler DJ. West Nile virus. *Lancet Infect Dis*. 2002; 2(9): 519-29.
5. Bunning ML, Bowen RA, Cropp CB, Sullivan KG, Davis BS, Komar N, et al. Experimental infection of horses with West Nile virus. *Emerg Infect Dis*. 2002; 8(4): 380-6.
6. Hayes EB, Sejvar JJ, Zaki SR, Lanciotti RS, Bode AV, Campbell GL. Virology, pathology, and clinical manifestations of West Nile virus disease. *Emerg Infect Dis*. 2005; 11(8): 1174-9.
7. Ostlund EN, Andresen JE, Andresen M. West Nile encephalitis. *Vet Clin North Am Equine Pract*. 2000; 16(3): 427-41.
8. Morales MA, Barrandeguy M, Fabbri C, Garcia JB, Vissani A, Troño K, et al. West Nile virus isolation from equines in Argentina. *Emerg Infect Dis*. 2006; 12(10): 1559-61.
9. Osorio JE, Ciuderis KA, Lopera JG, Piedrahita LD, Murphy D, LeVasseur J, et al. Characterization of West Nile viruses isolated from captive American flamingoes (*Phoenicopterus ruber*) in Medellín, Colombia. *Am J Trop Med Hyg*. 2012; 87(3): 565-72.
10. Pauvolid-Corrêa A, Morales MA, Levis S, Figueiredo LTM, Couto-Lima D, Campos Z, et al. Neutralising antibodies for West Nile virus in horses from Brazilian Pantanal. *Mem Inst Oswaldo Cruz*. 2011; 106(4): 467-74.

11. Melandri V, Guimarães AE, Komar N, Nogueira ML, Mondini A, Fernandez-Sesma A, et al. Serological detection of West Nile virus in horses and chicken from Pantanal, Brazil. *Mem Inst Oswaldo Cruz*. 2012; 107(8): 1073-5.
12. Silva JR, de Medeiros LC, dos Reis VP, Chávez JH, Munhoz TD, Borges GP, et al. Serologic survey of West Nile virus in horses from Central-West, Northeast and Southeast Brazil. *Mem Inst Oswaldo Cruz*. 2013; 108(7): 921-3.
13. Vieira M, Aguiar A, Borba A, Guimarães H, Eulálio K, Albuquerque-Neto L, et al. Letter to the editor West Nile Fever in Brazil: sporadic case, silent endemic disease or epidemic in its initial stages? *Rev Inst Med Trop Sao Paulo*. 2015; 57(3): 276.
14. Hall WC, Crowell TP, Watts DM, Barros VLR, Kruger H, Pinheiro F, et al. Demonstration of yellow fever and dengue antigens in formalin-fixed paraffin-embedded human liver by immunohistochemical analysis. *Am J Trop Med Hyg*. 1991; 45(4): 408-17.
15. Azevedo RSS, de Sousa JR, Araujo MTF, Martins Filho AJ, de Alcantara BN, Araujo FMC, et al. In situ immune response and mechanisms of cell damage in central nervous system of fatal cases microcephaly by Zika virus. *Sci Rep*. 2018; 8(1): 1-11.
16. Lanciotti RS, Kerst AJ, Nasci RS, Marvin S, Mitchell CJ, Savage HM, et al. Rapid detection of West Nile virus from human clinical specimens, field-collected mosquitoes, and avian samples by a TaqMan reverse transcriptase-PCR assay. *J Clin Microbiol*. 2000; 38(11): 4066-71.
17. Lanciotti RS, Kerst AMYJ. Nucleic acid sequence-based amplification assays for rapid detection of West Nile and St. Louis encephalitis viruses. *J Clin Microbiol*. 2001; 39(12): 4506-13.
18. Peng Y, Leung HCM, Yiu SM, Chin FYL. IDBA-UD: a *de novo* assembler for single-cell and metagenomic sequencing data with highly uneven depth. *Bioinformatics*. 2012; 28(11): 1420-8.
19. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol*. 2012; 19(5): 455-77.
20. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND. *Nat Methods*. 2015; 12(1): 59-60.
21. Katoh K, Standley DM. MAFFT. Multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol*. 2013; 30(4): 772-80.
22. Huson DH, Bryant D. Application of phylogenetic networks in evolutionary studies. *Mol Biol Evol*. 2006; 23(2): 254-67.
23. Posada D. jModelTest: phylogenetic model averaging. *Mol Biol Evol*. 2008; 25(7): 1253-6.
24. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics*. 2014; 30(9): 1312-3.
25. Rambaut A, Drummond AJ. Tracer V1.7.1. 2018. Available from <http://beast.bio.ed.ac.uk/Tracer>.
26. Rambaut A. FigTree, a graphical viewer of phylogenetic trees. Institute of Evolutionary Biology University of Edinburgh, 2009.
27. Deardorff E, Estrada-Franco J, Brault AC, Navarro-Lopez R, Campomanes-Cortes A, Paz-Ramirez P, et al. Introductions of West Nile virus strains to Mexico. *Emerg Infect Dis*. 2006; 12(2): 314-8.
28. Makino Y, Tadano M, Saito M, Fukunaga T, Maneekarn N, Sittisombut N, et al. Studies on serological cross-reaction in sequential flavivirus infections. *Microbiol Immunol*. 1994; 38(12): 951-5.
29. MS/SVS/DVDT - Ministério da Saúde/Secretaria de Vigilância em Saúde/Departamento de vigilância das Doenças Transmissíveis. Manual de vigilância sentinela de doenças neuroinvasivas por arbovírus. Brasília: Ministério da Saúde; 2017.
30. PAHO - Pan American Health Organization. Guía para la vigilancia, detección y respuesta para las encefalitis equinas. Washington (DC): OPS; 2013.